

# Precision-tinted spectral filters

A successful intervention for medically-resistant headaches and migraines after brain injury

SANDRA TOSTA, PHD AND HELEN IRLÉN, MFT  
JEFFREY LEWINE, PHD, JOSEPH ANNIBALI, MD



## Introduction

Precision-tinted spectral filters remedy perceptual difficulties and physical symptoms (including headaches and migraines) associated with the perceptual processing disorder, Irlen Syndrome. Advanced brain-mapping technology shows both anatomical and functional changes in brain activity that correspond directly with a reduction of symptoms when the correct color is worn as glasses. In March 2011, Irlen partnered with Semper Fi Fund to provide Irlen Spectral Filters to military personnel experiencing “medically resistant” headaches and migraines as a result of combat-related brain or head trauma.

## Objective

Extend the current therapeutic use of precision-tinted spectral filters to individuals who have acquired headaches and migraines as a result of head injury or brain trauma.

## Conclusion

For military personnel experiencing debilitating headaches and migraines that fail to respond to other interventions, Irlen Spectral Filters provide immediate relief and improvements. These improvements extend beyond headaches and migraines to impact other areas of life and daily functioning that are crucial to both their ability to remain in active duty and to achieve success after retiring from the military. Irlen Spectral Filters are able to eliminate the pain and discomfort that other interventions or medication have not been able to eliminate, provide stability and clarity in the visual field, and improve physical, emotional, and academic functioning. Future research should utilize brain imaging to document the reported improvements as functional and/or anatomical changes in the brain.

## References

Chouinard, B.D., Zhou, C.I., Hrybouski, S., Kim, E.S., Cummine, J. (2012). A functional neuroimaging case study of Meares-Irlen syndrome/visual stress (MISVIS) Brain Topography, 25 (3), 293-307.

Huang, J., Zong, X., Wilkins, A., Jenkins, B., Bozoki, A., Cao, Y. (2011). fMRI evidence that precision ophthalmic tints reduce cortical hyperactivation in migraine. Cephalgia, 31(8), 925-36.

Lewine, J.D., Irlen, H.L., & Orrison, W.W. (1996). Visual evoked magnetic fields in Scotopic sensitivity syndrome. (Available from New Mexico Institute of Neuroimaging. The New Mexico Regional Field Medical Center: Albuquerque, NM).

Riddell, P. M., Wilkins, A. and Hainline, L. (2006). The effect of colored lenses on the visual evoked response in children with visual stress. Optometry and Vision Science, 83 (5), 299-305.

## Acknowledgements

This work was supported in part by the Semper Fi Foundation. Special thanks to Mauricio Ferreira, PhD, Eugene Oh, and the Hypothesis Group for their analytic contribution.

## For More Info

Sandra Tosta  
sandy@irlen.com  
www.irlen.com  
5380 Village Rd.,  
Long Beach, CA 90808  
562-496-2550

## Methods


### 178 U.S. military personnel

Have already tried one or more treatments: Migraine medication, other medication, acupuncture, chiropractic treatments, healing touch, yoga, meditation, vestibular therapy, botox, hyperbaric, alpha stem

Data regarding headache and migraine frequency and medication use were analyzed using paired-samples t-tests. All comparisons were highly significant ( $p=.00$ ). Likert scale data related to physical symptoms and activities causing difficulties were analyzed using multiple analysis of variance (MANOVA) with Bonferroni correction. The multivariate and all univariate analyses were highly significant ( $p=.00$ ).

### Areas of inquiry

Frequency/severity of headaches/migraines, light sensitivity, academic difficulties, physical symptoms, emotional symptoms, depth perception, driving



## Results



### Percent of the day headaches are experienced (avg)



### SEVERITY OF DIFFICULTY BEFORE AND AFTER

Average rating of difficulty (0-5 scale, 0 means “no problem” and 5 means “severe problem”)

